

NASA-CR-193503

WORKSHOP ON EARLY MARS: HOW WARM AND HOW WET?


0194104
14 P

(NASA-CR-193503) WORKSHOP ON EARLY
MARS: HOW WARM AND HOW WET, PART 2?
(Lunar and Planetary Inst.) 14 p

N94-17637

Unclass

G3/91 0194104



MSATT

Mars Surface and Atmosphere Through Time



LPI Technical Report Number 93-03, Part 2

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058-1113
LPI/TR--93-03, Part 2



WORKSHOP ON
EARLY MARS: HOW WARM AND HOW WET?

Edited by
S. Squyres and J. Kasting

Held in
Breckenridge, Colorado

July 26–28, 1993

Sponsored by
MSATT Study Group
Lunar and Planetary Institute

Lunar and Planetary Institute 3600 Bay Area Boulevard Houston TX 77058-1113

LPI Technical Report Number 93-03, Part 2
LPI/TR--93-03, Part 2

Compiled in 1993 by
LUNAR AND PLANETARY INSTITUTE

The Institute is operated by the University Space Research Association under Contract No. NASW-4574 with the National Aeronautics and Space Administration.

Material in this volume may be copied without restraint for library, abstract service, education, or personal research purposes; however, republication of any paper or portion thereof requires the written permission of the authors as well as the appropriate acknowledgment of this publication.

This report may be cited as

Squyres S. and Kasting J., eds. (1993) *Workshop on Early Mars: How Warm and How Wet?* LPI Tech. Rpt. 93-03, Part 2, Lunar and Planetary Institute, Houston. 23 pp.

This report is distributed by

ORDER DEPARTMENT
Lunar and Planetary Institute
3600 Bay Area Boulevard
Houston TX 77058-1113

Mail order requestors will be invoiced for the cost of shipping and handling.

Preface

In 1992 the MSATT program conducted a workshop on modeling of the martian climate. At that workshop it became clear that a serious problem had arisen concerning the early climate of Mars. Based on the evidence for small-scale fluvial activity, the view had been widely held that early in its history Mars had a climate that was much warmer and wetter than today's. However, most plausible recent climate models have fallen far short of the warm temperatures often inferred from the geologic evidence. Moreover, recent geophysical work has suggested that early geothermal warming may also have played a significant role in allowing fluvial activity. In order to address the issue of just how warm and how wet early Mars was, a workshop was convened in July of 1993, in Breckenridge, Colorado. The results of the workshop are reported here.

PRECEDING PAGE BLANK NOT FILMED

Contents

Summary of Technical Sessions	1
List of Workshop Participants	3

PRECEDING PAGE BLANK NOT FILMED

Program

Monday, July 26, 1993

8:00 a.m. *Registration*

8:45–9:45 a.m.

OVERVIEW OF MARS' EARLY EVOLUTION

8:45 a.m. *Early Mars: Warmed from Without or Within*
S. W. Squyres*

9:15 a.m. *Possible Solutions to the Problem of Channel Formation on Early Mars*
J. F. Kasting*

9:45 a.m. *Break*

10:00 a.m.–12:30 p.m.

VOLATILE INVENTORIES AND ATMOSPHERIC EVOLUTION

10:00 a.m. *Wet Inside and Out?: Constraints on Water in the Martian Mantle and on Outgassed Water, Based on Melt Inclusions in SNC Meteorites*
H. Y. McSween Jr.* and R. P. Harvey

10:30 a.m. *Mars Atmospheric Loss and Isotopic Fractionation by Solar-Wind-induced Sputtering and Photochemical Escape*
B. M. Jakosky*, R. O. Pepin, R. E. Johnson, and J. L. Fox

11:00 a.m. *Evolution of the Martian Atmosphere*
R. O. Pepin*

11:30 a.m. *Requirements for the Early Atmosphere of Mars from the Nitrogen Isotope Ratios*
J. L. Fox*

12:00 p.m. *Adjourn*

7:00–9:00 p.m.

CLIMATE/GEOCHEMICAL/PHOTOCHEMICAL MODELING

7:00 p.m. *A Model for the Evolution of CO₂ on Mars*
R. M. Haberle*, D. Tyler, C. P. McKay, and W. L. Davis

* Indicates speaker

- 7:30 p.m. *Early Mars: The Inextricable Link Between Internal and External Influences on Valley Network Formation*
S. E. Postawko* and F. P. Fanale
- 8:00 p.m. *The Young Sun and Photochemistry of the Primitive Martian Atmosphere*
H. Nair*, M. F. Gerstell, and Y. L. Yung
- 8:30 p.m. *A Carbon Dioxide/Methane Greenhouse Atmosphere on Early Mars*
L. L. Brown* and J. F. Kasting
- 9:00 p.m. *Mars and the Early Sun*
D. Whitmire*, L. R. Doyle, R. T. Reynolds, and P. G. Whitman
- 9:30 p.m. *Adjourn*

Tuesday, July 27, 1993

9:00 a.m.–12:30 p.m.

MARTIAN HYDROLOGY AND VALLEY SYSTEM FORMATION

- 9:00 a.m. *The Changes on Mars at the End of Heavy Bombardment*
M. H. Carr*
- 9:30 a.m. *Evolution of the Global Water Cycle on Mars: The Geological Evidence*
V. R. Baker* and V. C. Gulick
- 10:00 a.m. *Fluvial Valleys in the Heavily Cratered Terrains of Mars: Evidence for Paleoclimatic Change?*
V. C. Gulick* and V. R. Baker
- 10:30 a.m. *Break*
- 11:00 a.m. *The Hydrologic Response of Mars to the Onset of a Colder Climate and to the Thermal Evolution of Its Early Crust*
S. M. Clifford*
- 11:30 a.m. *Briny Lakes on Early Mars? Terrestrial Intracrater Playas and Martian Candidates*
P. Lee*
- 12:00 p.m. *Adjourn*

7:00–9:30 p.m.

CLUES FROM MARTIAN GEOMORPHOLOGY

- 7:00 p.m. *Early Mars: A Regional Assessment of Denudation Chronology*
T. A. Maxwell* and R. A. Craddock
- 7:30 p.m. *The Early Martian Environment: Clues from the Cratered Highlands and the Precambrian Earth*
R. A. Craddock* and T. A. Maxwell
- 8:00 p.m. *Ancient Martian Valley Genesis and Paleoclimatic Inference: The Present as a Key to the Past*
G. R. Brakenridge*
- 8:30 p.m. *Mars: Noachian Hydrology by Its Statistics and Topology*
N. A. Cabrol* and E. A. Grin
- 9:00 p.m. *The Martian Valley Networks: Origin By Niveo-Fluvial Processes*
J. Rice*
- 9:30 p.m. *Adjourn*

Wednesday, July 28, 1993

9:00—10:00 a.m.

CHEMICAL WEATHERING AND CLIMATE

- 9:00 a.m. *An Attempt to Comprehend Martian Weathering Conditions Through the Analysis of Terrestrial Palagonite Samples*
C. Douglas*, I. P. Wright, J. B. Bell, R. V. Morris, D. C. Golden,
and C. T. Pillinger
- 9:30 a.m. *Oxidation of Dissolved Iron Under Warmer, Wetter Conditions on Mars: Transitions to Present-Day Arid Environments*
R. G. Burns*
- 10:00 a.m. *Break*
- 10:30 a.m. *Discussion*
- 12:30 p.m. *Adjourn*

Summary of Technical Sessions

It is clear that conditions on Mars during the Noachian Period (which extended from the beginning of Mars' geologic record to something like 3.9 b.y. ago) were significantly different than they are today. As on the other planets, the rate of meteoritic bombardment was far higher than it is at present; this is shown clearly by the densely cratered nature of Noachian terrains. But there were other differences as well. Some are only inferred, while for others we have clear evidence. The nature of these differences, and what they imply about broader topics of planetary evolution, were the primary subjects of the workshop.

The atmosphere of Mars during the Noachian was probably substantially more dense than the present atmosphere of 7 mbar of CO₂. Just how dense it was is a subject of some debate. Processes that fed it during this period include volcanic outgassing and infall of volatile-rich meteorites. Processes that destroyed it may have included hydrodynamic and thermal escape, solar wind erosion, impact erosion, and chemical processes at the surface such as carbonate formation. The balance among these various processes has been a topic of much recent work and was vigorously discussed at the workshop. While the atmospheric pressure during the Noachian is unknown, there is a consensus that pressures of up to a few bars are plausible. If this was the case, then carbonate formation during or after the Noachian must have been important. Other sinks could remove hundreds of millibars of CO₂ from the atmosphere, but removal of a few bars would require that a substantial carbonate reservoir currently be present on Mars.

Probably another important characteristic of the Noachian was a much higher geothermal heat flow. Detailed models of the accretion and early thermal evolution of Mars, as well as isotopic evidence from SNC meteorites believed to come from Mars, provide persuasive evidence for very early core formation, a hot early mantle, and a high early heat flow. This recognition that Mars probably began its history hot has been one of the major developments in martian geophysics over the last several years. The mean global heat flow during the Noachian is also very uncertain, but a value 3.9 b.y. ago of about 5 × the present one (which is typically estimated at about 45–50 mW/m²) is plausible. Whatever the global mean heat flow was at that time, it is certain that local values in areas of enhanced geothermal activity were significantly higher still.

Given these very different environmental conditions, it is natural to expect that geomorphic processes during the Noachian were different as well, and it is clear that they were. The most obvious difference is that the rate of formation of valley systems was far higher during the Noachian than it has

been during any other period of martian history. Valley systems are small, quasidendritic drainage networks found primarily in the ancient cratered highlands of Mars. Typical valley widths are 1–2 km, and lengths extend up to several hundred kilometers. The valleys were almost certainly carved by liquid water flowing at low discharges in small channels. These valley systems have long provided the primary physical evidence suggesting that conditions on Mars were different early in the planet's history.

There is other morphologic evidence as well. Careful comparison of the depths and rim heights of Noachian craters with post-Noachian craters shows clearly that the rate of crater degradation (lowering of crater rims and raising of crater floors) was far higher in the Noachian than it has been since then. Such degradation is certainly attributable in part to erosion and deposition, though other processes (such as volcanic infilling) could also have played a role.

There is general agreement that there has been enough H₂O on Mars for us to accept that water was the fluid that carved the valley systems. However, the details of their formation are unclear. In particular, there are two key obviously linked questions that remain unresolved: Did conditions during the Noachian allow precipitation? And does the geomorphic evidence from the Noachian require precipitation?

If the climate of Mars was warm enough during the Noachian to allow precipitation, then an enhanced greenhouse effect enabled by a denser atmosphere is required. However, temperatures anywhere close to the melting point of water may be difficult to achieve with a dense early CO₂ atmosphere on Mars. Earlier models predicted substantial greenhouse warming for an atmosphere of a few bars of CO₂. However, more recent models that include the effects of CO₂ condensation and cloud formation suggest that such an atmosphere is capable of raising the martian surface temperature by at most several tens of degrees. For the conventional view of solar evolution (which suggests a luminosity about 75% the present value during the Noachian), such a greenhouse does not appear capable of producing mean surface temperatures in excess of ~230 K.

There are two other factors that could conceivably contribute further climatic warming, but both are highly uncertain. One is the possibility that the early Sun was substantially more luminous than predicted by conventional models; this would require a much higher rate of solar mass loss during the Sun's early history than is usually accepted. The other is the possibility that substantial quantities of methane, another very effective greenhouse gas, were present in the martian

atmosphere. Without either of these effects, however, the best recent work suggests that climatic warming to temperatures above 273 K during the Noachian was not possible.

There also is no general agreement as to whether the geomorphic evidence actually requires precipitation. Some researchers point to the ubiquitous nature of the valley systems in old heavily cratered terrain on Mars, and to the fact that some valley heads extend right to the crests of drainage divides. From these characteristics they infer that precipitation and runoff must have taken place. Others note that the morphology of many valley systems is more consistent with sapping than runoff, and that the highest drainage densities on Mars are substantially lower than those in even arid regions on Earth. From these characteristics, they infer that the valleys formed by sapping of subsurface liquid, and that precipitation was not required. There is also disagreement as to whether the early high rates of crater degradation require precipitation, or whether they are just a consequence of higher early rates of sapping, eolian transport, volcanism, and so forth.

There are geomorphic points on which there is agreement. There are some valley systems for which all workers agree that sapping is a satisfactory explanation, without the need to invoke precipitation and runoff. It is also clear that many valley systems are too extensive to have formed by the single discharge of an aquifer, i.e., that some type of aquifer recharge is required. Several different processes, including precipitation and hydrothermal convection (enabled by the early high heat flow), have been examined closely as recharge mechanisms. Both of these appear to be more than adequate to do the job. The basic difficulties are that dissimilar processes can produce similar final landforms, and that many

processes may have operated on these valleys, including ones that have substantially obscured the evidence for the primary genetic process.

A final issue that was considered at the workshop was identification of directions for future work. Important areas where further progress should be possible include these:

- Use geochemical data from Mars Observer or a follow-on mission to search for martian carbonates.
- Use imaging data from Mars Observer or a follow-on mission to search for further morphologic evidence requiring Noachian precipitation.
- Improve models of the contribution of methane to the martian greenhouse.
- Further investigate the prospects for a more luminous early Sun.
- Model the climatic effects of the very high and rapidly varying obliquities that are suggested by recent dynamical modeling.
- Improve climate models to include the optical properties of CO₂ clouds.
- Continue investigation of terrestrial analog terrains and processes.

Note added in proof: In light of the failure of Mars Observer, a top priority should be a reflight of the instruments to obtain the data that otherwise will not be acquired.

List of Workshop Participants

J. R. Baker

*Department of Geosciences
University of Arizona
Tucson AZ 85721
Phone: 602-621-6003
Fax: 602-621-2672*

J. Bergstralh

*Code SLC
NASA Headquarters
Washington DC 20546
Phone: 202-453-1580
Fax: 202-358-3097
E-mail: NHQVAX::JBERGSTR*

Robert Brakenridge

*Surficial Processes Laboratory
Department of Geography
Dartmouth College
Hanover NH 03755
Phone: 603-646-2870
Fax: 603-646-2810
E-mail: brakenridge@dartmouth.edu*

Lisa L. Brown

*Department of Geosciences
332 Deike Building
Pennsylvania State University
University Park PA 16802
Phone: 814-863-3965
Fax: 814-865-3191
E-mail: brown@essc.psu.edu*

Mark Bullock

*Laboratory for Atmospheric and Space Physics
Campus Box 392
University of Colorado
Boulder CO 80309-0392
Phone: 303-492-1628
Fax: 303-492-6946
E-mail: ZODIAC::BULLOCK*

Roger G. Burns

*Mail Code 54-816
Massachusetts Institute of Technology
Cambridge MA 02139
Phone: 617-253-1906
Fax: 617-253-6208
E-mail: bcabral@eagle.mit.edu*

Bryan Butler

*Division of Geological and Planetary Science
Mail Stop 170-25
California Institute of Technology
Pasadena CA 91125
Phone: 818-356-6477
Fax: 818-585-1917
E-mail: butler@scat.gps.caltech.edu*

Nathalie Cabrol

*Telescope de 1 Metre
Observatoire de Paris
92195 Meudon Cedex Principal
FRANCE
Phone: 45-07-75-30
Fax: 45-07-79-59
E-mail: MESIOA::DOLLFUS*

Michael H. Carr

*Mail Stop 946
U.S. Geological Survey
345 Middlefield Road
Menlo Park CA 94025
Phone: 415-329-5174
Fax: 415-329-4936
E-mail: mcarr@isdmnl.wr.usgs.gov*

Michael Carroll

*Sky and Telescope Magazine
6820 W. Chestnut Avenue
Littleton CO 80123
Phone: 303-933-1645*

Benton Clark

*Mail Stop B0560
Martin Marietta
P.O. Box 179
Denver CO 80201
Phone: 303-971-9007
Fax: 303-971-9141
E-mail: bclark@den.munc.com*

Stephen Clifford

*Lunar and Planetary Institute
3600 Bay Area Boulevard
Houston TX 77058
Phone: 713-486-2146
Fax: 713-486-2162
E-mail: clifford@lpi.jsc.nasa.gov*

Robert A. Craddock

*Center for Earth and Planetary Studies
National Air and Space Museum, Room 3775
MRC 315
Smithsonian Institution
Washington DC 20560
Phone: 202-357-1457
Fax: 202-786-2566
E-mail: craddock@ceps.edu*

David DesMarais

*Mail Stop 239-4
NASA Ames Research Center
Moffett Field CA 94035-1000
Phone: 415-604-3220
Fax: 415-604-1088
E-mail: david_desmarais@qmgate.arc.nasa.gov*

Tammy Dickinson

*Code SLC
NASA Headquarters
Washington DC 20546-0001
Phone: 202-358-0292
Fax: 202-358-3097
E-mail: tdickinson@nasamail.nasa.gov*

Caroline Douglas

*Planetary Science Unit
Department of Earth Sciences
Walton Hall
Open University
Milton Keynes MK7 6AA
UNITED KINGDOM
Phone: 0908-655-988
Fax: 0908-655-151*

Gerlind Dreibus

*Max-Planck-Institut für Chemie
Saarstrasse 23
D-6500 Mainz
GERMANY
Phone: 49-6131-305-395
Fax: 49-6131-371-290*

Jane L. Fox

*MSRC
Institute for Terrestrial and Planetary Atmospheres
State University of New York
Stony Brook NY 11794
Phone: 516-632-8317
Fax: 516-632-8820
E-mail: fox@atmsci.sunysb.edu*

Virginia Gulick

*Mail Stop 245-3
Space Science Division
NASA Ames Research Center
Moffett Field CA 94035
Phone: 415-604-0781
Fax: 415-604-6779
E-mail: gulick@convx1.ccit.arizona.edu*

R. M. Haberle

*Mail Stop 245-3
Space Science Division
NASA Ames Research Center
Moffett Field CA 94035
Phone: 415-604-5491
Fax: 415-604-6779
E-mail: haberle@humbabe.arc.nasa.gov*

Joan Hayashi

*Laboratory for Atmospheric and Space Physics
Campus Box 392
University of Colorado
Boulder CO 80309
Phone: 303-492-8918
Fax: 303-492-6946
E-mail: hayashi@argyre.colorado.edu*

Bruce Jakosky

*Laboratory for Atmospheric and Space Physics
Campus Box 392
University of Colorado
Boulder CO 80309-0392
Phone: 303-492-8004
Fax: 303-492-6946
E-mail: ZODIAC::JAKOSKY*

John Jones

*Mail Code SN2
NASA Johnson Space Center
Houston TX 77058
Phone: 713-483-5319
Fax: 713-483-5347
E-mail: SN::JJONES*

James Kasting

*211 Deike
Pennsylvania State University
University Park PA 16802
Phone: 814-865-3207
Fax: 814-865-3191
E-mail: kasting@essc.psu.edu*

Pascal Lee

404 Space Sciences Building
 Cornell University
 Ithaca NY 14853-6801
 Phone: 607-255-6237
 Fax: 607-255-9002
 E-mail: lee@astronsun.tn.cornell.edu

Ted A. Maxwell

National Air and Space Museum
 MRC 315
 Smithsonian Institution
 Washington DC 20560
 Phone: 202-357-1424
 Fax: 202-786-2566
 E-mail: tmaxwell@nasamail.nasa.gov

Daniel J. McCleese

Mail Stop 183-335
 Jet Propulsion Laboratory
 4800 Oak Grove Drive
 Pasadena CA 91109
 Phone: 818-354-2317
 Fax: 818-393-6546
 E-mail: djmcc@scn1.jpl.nasa.gov

Harry McSween

Department of Geological Sciences
 University of Tennessee
 Knoxville TN 37996-1410
 Phone: 615-974-5498
 Fax: 615-974-2368
 E-mail: mcsween@utkvtx.utk.edu

Mike Mellon

Laboratory for Atmospheric and Space Physics
 Campus Box 392
 University of Colorado
 Boulder CO 80302
 Phone: 303-492-7902
 Fax: 303-492-6946
 E-mail: mellon@argyre.colorado.edu

Michael A. Meyer

Lockheed Engineering and Sciences Co.
 500 E Street, SW
 Suite 800
 Washington DC 20024
 Phone: 202-863-5257
 Fax: 202-863-8217
 E-mail: mmeyer@nasamail.nasa.gov

Hari A. Nair

Mail Stop 170-25
 California Institute of Technology
 Pasadena CA 91125
 Phone: 818-356-6960
 Fax: 818-585-1917
 E-mail: hari@mercul.gps.caltech.edu

Robert O. Pepin

School of Physics and Astronomy
 University of Minnesota
 116 Church Street, SE
 Minneapolis MN 55455
 Phone: 612-624-0819
 Fax: 612-624-4578

Jeff Plescia

Mail Stop 183-501
 Jet Propulsion Laboratory
 4800 Oak Grove Drive
 Pasadena CA 91109
 Phone: 818-354-6936
 Fax: 818-354-0966
 E-mail: jplescia@nasamail.nasa.gov

Susan Postawko

School of Meteorology
 University of Oklahoma
 100 East Boyd
 Norman OK 73019
 Phone: 405-325-6561
 Fax: 405-325-7689
 E-mail: spostawko@geohub.gen.uoknor.edu

J. Rice

Department of Geography
 Arizona State University
 Tempe AZ 85287
 Phone: 602-496-6512
 Fax: 602-965-8313
 E-mail: asjwr@asuvm.inre.asu.edu

Ben Schuraytz

Lunar and Planetary Institute
 3600 Bay Area Boulevard
 Houston TX 77058
 Phone: 713-486-2187
 Fax: 713-486-2162
 E-mail: schuraytz@lpi.jsc.nasa.gov

Bernhard Spettel

*Abteilung Kosmochemie
Max-Planck-Institut für Chemie
Saarstrasse 23
D-6500 Mainz
GERMANY
Phone: 49-6131-305-233*

Robert Zubrin

*Martin Marietta
P.O. Box 179
Denver CO 80201
Phone: 303-971-9299
Fax: 303-971-3600*

Steve Squyres

*Space Sciences Building
Cornell University
Ithaca NY 14853
Phone: 607-255-3508
Fax: 607-255-5907
E-mail: squyres@astrosun.tn.cornell.edu*

John Stansberry

*Department of Planetary Sciences
University of Arizona
Tucson AZ 85721
Phone: 602-621-2276
E-mail: stansber@lpl.arizona.edu*

Dan Tyler Jr.

*2561 Mardell Way
Mountain View CA 94043
Phone: 415-604-0322
E-mail: tyler@oskar.ames.nasa.gov*

Daniel P. Whitmire

*Department of Physics 44210
University of Southwestern Louisiana
Lafayette LA 70504
Phone: 318-231-6185
Fax: 318-321-6190
E-mail: dpw9254@usl.edu*

Frank Wlotzka

*Abteilung Kosmochemie
Max-Planck-Institut für Chemie
Saarstrasse 23
D-6500 Mainz
GERMANY
Phone: 49-6131-305-343
Fax: 49-6131-371-290
E-mail: wlotzka@mpch-mainz.mpg.dbp.de*

Aaron Zent

*Mail Stop 245-3
NASA Ames Research Center
Moffett Field CA 94035
Phone: 415-604-5517
Fax: 415-604-6779
E-mail: zent@pan.arc.nasa.gov*